


* Cable Equation

①



r_i : intracellular axial resistance
 per unit length (Ω/m)
 r_o : extracellular " (Ω/m)
 times unit length
 r_m : membrane resistance ($\Omega \cdot m$)
 in radial direction
 C_m : membrane capacitance per unit
 length (F/m)

I_i : total longitudinal intracellular
 current (A)

I_o : " " extracellular
 (A)

i_m : total transmembrane current
 per unit length (A/m)

$i_{m,c}$: capacitive component of i_m

$i_{m,i}$: ionic "

(2)

Φ_i : potential inside membrane [V]

Φ_o : " outside " [V]

$V_m = \Phi_i - \Phi_o$: membrane voltage [V]

V_r : resting membrane voltage [V]

$V' = V_m - V_r$: deviation of membrane voltage from V_r

Without stimulating current, $I_i + I_o = 0$

At resting state, $I_i = I_o = I_m = 0$, $V_m = V_r$, $V' = 0$

During activation, $I_i + I_o = 0$, $V' \neq 0$

Since V_r is constant for all x ,

$$\frac{\partial V'}{\partial x} = \frac{\partial V_m}{\partial x}, \quad \frac{\partial V'}{\partial t} = \frac{\partial V_m}{\partial t}$$

② In steady state, $\frac{\partial}{\partial t} = 0$.

$$\frac{\partial \Phi_i}{\partial x} = -I_i r_i, \quad \frac{\partial \Phi_o}{\partial x} = -I_o r_o$$

$$\dot{\lambda}_m = -\frac{\partial I_i}{\partial x} = \frac{\partial I_o}{\partial x}$$

(loss of I_i) (gain of I_o)

$$\frac{\partial V'}{\partial x} = \frac{\partial \Phi_i}{\partial x} - \frac{\partial \Phi_o}{\partial x} = -I_i r_i + I_o r_o \quad (3)$$

$$\frac{\partial^2 V'}{\partial x^2} = -r_i \underbrace{\frac{\partial I_i}{\partial x}}_{-i_m} + r_o \underbrace{\frac{\partial I_o}{\partial x}}_{i_m} = (r_i + r_o) i_m$$


Since $C_m \frac{dV'_o}{dt} = 0$, $i_m = \frac{V'}{r_m}$

$$\frac{\partial^2 V'}{\partial x^2} = \frac{r_i + r_o}{r_m} V'$$

$$\frac{\partial^2 V'}{\partial x^2} - \lambda^2 V' = 0, \quad \lambda = \sqrt{\frac{r_m}{r_i + r_o}} \approx \sqrt{\frac{r_m}{r_i}}$$

$$V'_{x=0} = V'(0), \quad V'_{x=\infty} = 0$$

$$V'(x) = V'(0) e^{-\frac{x}{\lambda}}$$



④

② Step-current stimulation

$$\dot{I}_m = \dot{I}_{m_i} + \dot{I}_{m_e}$$

$$\frac{I}{r_i + r_o} \frac{\partial^2 V'}{\partial x^2} = \frac{V'}{r_m} + C_m \frac{\partial V'}{\partial t}$$

$$\frac{r_m}{r_i + r_o} \frac{\partial^2 V'}{\partial x^2} - V' - r_m C_m \frac{\partial V'}{\partial t} = 0$$

$$-\lambda^2 \frac{\partial^2 V'}{\partial x^2} + \tau \frac{\partial V'}{\partial t} + V' = 0$$